arXiv:1604.06794v1 [math.HO] 21 Apr 2016

CYCLIC EXTENSIONS ARE RADICAL

MARIANO SUÁREZ-ÁLVAREZ

The fact that finite Galois extensions with cyclic Galois group can be constructed by adjoining a root of a well-chosen element of the base field is a basic result of Galois theory and a key point in studying the problem of solvability by radicals. In textbooks on the subject, this result is usually obtained as a consequence of Hilbert's Theorem 90, which is itself deduced from Artin's theorem on the independence of characters. The very simple argument we present below sidesteps those requirements.

Theorem. *If E/K is a cyclic extension of degree n of a field K which contains a primitive nth root of unity ζ, then there exists an $x \in E$ with $x^n \in K$ and $E = K(x)$.*

Notice that the existence of a primitive nth root of unity in K implies that the characteristic of this field does not divide n.

Proof. Let σ be a generator of the Galois group G of the extension. Since $\sigma^n = \mathrm{id}_E$, the minimal polynomial of σ over K divides $X^n - 1$ in $K[X]$ and then, as this polynomial has all its roots in K and they are all simple, σ is diagonalizable and its eigenvalues are nth roots of unity.

Let Γ be the set of eigenvalues of σ. If λ, $\mu \in \Gamma$, so that there are a, $b \in E^\times$ such that $\sigma(a) = \lambda a$ and $\sigma(b) = \mu b$, then $\lambda\mu \in \Gamma$, as $\sigma(ab) = \lambda\mu ab$. Since Γ is contained in Ω_n, the finite group of nth roots of unity, this is enough to conclude that Γ is in fact a *subgroup* of Ω_n. If m is the order of Γ , then we have that $\lambda^m = 1$ for all $\lambda \in \Gamma$ and, as σ is diagonalizable with eigenvalues in Γ, that $\sigma^m = \mathrm{id}_E$. As the order of σ is n and $m \leq n$, it follows from this that $m = n$.

All nth roots of unity are therefore eigenvalues of σ and, in particular, there is an $x \in E^\times$ such that $\sigma(x) = \zeta x$. Then $\sigma(x^n) = \zeta^n x^n = x^n$, so x^n is in the fixed field $E^G = K$. We thus see that x is a root of the polynomial $X^n - x^n$ of $K[X]$, which is irreducible over K: indeed, it has simple roots x, ζx, $\ldots$, $\zeta^{n-1}x$ in E and these are permuted transitively by σ. The degree of the subextension $K(x)/K$ of E/K is then n and, of course, this means that $E = K(x)$. □

E-mail address: mariano@dm.uba.ar

Departamento de Matemática. Facultad de Ciencias Exactas y Naturales. Universidad de Buenos Aires. Ciudad Universitaria, Pabellón I. (1428) Ciudad de Buenos Aires (Argentina)

Date: November 8, 2014.

2010 *Mathematics Subject Classification.* 12F05.

This work was partially supported by UBACYT X475, PIP-CONICET 2012-2014 11220110100870, PICT 2011-1510 and MathAmSud-GR2HOPF..